


2022
PEDIATRIC
MEASURES
GUIDE



Hackensack Meridian
Partners

Summary HMHP Contract Measures	3
Appropriate Testing for Pharyngitis (CWP)	4
Asthma Medication Ratio (AMR)	5
Child And Adolescent Well-Care Visits (WCV)	6
Depression Screening - 12 Years and Older	7
Weight Assessment and Counseling for Nutrition and Physical Activity For Children/Adolescents (WCC)	8
Well-Child Visits in the First 30 Months of Life (W30)	9



	LINE OF BUSINESS:	MEDICARE			COMMERCIAL			HMHP
CATEGORY	PAYOR:	Aetna	Braven	Horizon	Aetna	Cigna	Horizon	HMHP
Access	Child and Adolescent Well Care - 3 to 21 years old					X		X
	Well-Child Visits in the First 30 Months of Life (Two sub-measures: 0-15 months is 6 or more visits; 15-30 months is 2 or more visits)							X
Chronic Disease Management	Depression Screening					X		
Medication Management	Appropriate Testing for Pharyngitis							X
	Asthma Medication Ratio - 5 to 64 years						X	X
Preventive Care Screenings	Weight Assessment and Counseling for Nutrition and Physical Activity for Childhood/Adolescents – BMI Percentile							X
	Weight Assessment and Counseling for Nutrition and Physical Activity for Childhood/Adolescents – Counseling for Nutrition							X

1 CPT® is a registered trademark of the American Medical Association. ICD-10-CM is the International Classification of Diseases, Tenth Revision, Clinical Modification. Source: MA 5 Star Technical Specifications and NCQA HEDIS® 2022; Technical Specifications for Health Plans HEDIS® measures and specifications are not a substitute for clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications. Documentation is for training and educational purposes.

APPROPRIATE TESTING FOR PHARYNGITIS (CWP)

<p>Measure Description</p> <p>Measure Source: HEDIS®</p>	<p>The percentage of episodes for patients 3 years of age and older where the patient was diagnosed with pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test in the seven-day period from three days prior to the Episode Date through three days after the Episode Date. A higher rate represents better performance (i.e., appropriate testing). Three age ranges are reported: 3 -17 years; 18-64 years; 65 and older; and a total.</p>
<p>Exclusions</p>	<p>Patients who have Negative Medication History:</p> <ul style="list-style-type: none"> ■ A period of 30 days prior to the Episode Date when the member had no pharmacy claims for either new or refill prescriptions for a listed antibiotic drug. ■ No prescriptions filled more than 30 days prior to the Episode Date that are active on the Episode Date. <p>Patients who have a Comorbid Condition History:</p> <ul style="list-style-type: none"> ■ HIV, Malignant Neoplasms, Emphysema, COPD, and immune disorders. <p>Patients who are in hospice are excluded from the measure.</p>
<p>Medications</p>	<p>Amoxicillin, Ampicillin, Amoxicillin-clavulanate, Cefadroxil, Cefazolin, Cephalexin, Trimethoprim, Clindamycin, Azithromycin, Clarithromycin, Erythromycin, Erythromycin ethylsuccinate, Erythromycin lactobionate, Erythromycin stearate, Penicillin G potassium, Penicillin G sodium, Penicillin V potassium, Penicillin G Benzathine, Dicloxacillin, Ciprofloxacin, Levofloxacin, Moxifloxacin, Ofloxacin, Cefaclor, Cefprozil, Cefuroxime, Sulfamethoxazole-trimethoprim, Doxycycline, Minocycline, Tetracycline, Cefdinir, Cefixime, Cefpodoxime, Ceftibuten, Cefditoren, Ceftriaxone</p>
<p>Documentation</p>	<p>Claims based measure only. A group A streptococcus test in the seven-day period from three days prior to the IESD through three days after the IESD makes patients compliant.</p>
<p>Considerations</p>	<p>The denominator for this measure is based on episodes, not patients. All eligible episodes that were not excluded or deduplicated remain in the denominator.</p> <p>Only patients who were prescribed antibiotics and have a diagnosis are included in the denominator; if patients do not have a fill for the antibiotic, they will not be included in the measure.</p> <p>Patients seen for telehealth visits are included in the measure.</p>
<p>CODES:</p>	
<p>Pharyngitis ICD10</p>	<p>J02.0, J02.8-9, J03.00-.01, J03.80-.81, J03.90-9.1</p>
<p>Group A Strep Tests CPT</p> <p>LOINC</p> <p>SNOMED</p>	<p>87070, 87071, 87081, 87430, 87650-87652, 87880</p> <p>11268-0, 17656-0, 17898-8, 18481-2, 31971-5, 49610-9, 5036-9, 60489-2, 626-2, 6557-3 6558-1, 6559-9, 68954-7, 78012-2</p> <p>122121004, 122205003, 122303007</p>

1 CPT® is a registered trademark of the American Medical Association. ICD-10-CM is the International Classification of Diseases, Tenth Revision, Clinical Modification. Source: MA 5 Star Technical Specifications and NCQA HEDIS® 2022; Technical Specifications for Health Plans HEDIS® measures and specifications are not a substitute for clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications. Documentation is for training and educational purposes.

<p>Measure Description</p> <p>Measure Source: HEDIS®</p>	<p>The percentage of members 5-64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.</p> <p>Report the following age stratifications and total rate: ■ 5-11 years ■ 12-18 years ■ 19-50 years ■ 51-64 years ■ Total Rate</p> <p>Members are identified as having persistent asthma: ■ At least one ED visit, with a principal diagnosis of asthma. ■ At least one acute inpatient encounter, with a principal diagnosis of asthma without telehealth. ■ At least one acute patient discharge with a principal diagnosis of asthma. ■ At least four outpatient visits or observation visits, telephone visits, or online assessments, on different dates of service, with any diagnosis of asthma and at least two asthma medication dispensing events for any controller medication or reliever medication. Visit type need not be the same for the four visits. Only three of the four visits may be a telehealth visit, telephone visit, or online assessment. ■ At least four asthma medication dispensing events.</p>
<p>Exclusions</p>	<p>Members who had no asthma medications dispensed during the measurement year.</p> <ul style="list-style-type: none"> Members in hospice. Members with emphysema, COPD, obstructive chronic bronchitis, chronic respiratory conditions related to fumes/vapors, cystic fibrosis, and acute respiratory failure.
<p>Medications</p>	<p>Antiasthmatic combinations: Dyphylline-guaifenesin Antibody inhibitors: Omalizumab Anti-interleukin-4: Dupilumab Anti-interleukin-5: Benralizumab, Mepolizumab, Reslizumab Inhaled corticosteroids: Beclomethasone, Budesonide, Ciclesonide, Flunisolide, Fluticasone, Mometasone Inhaled steroid combinations: Budesonide-formoterol, Fluticasone-vilanterol, Fluticasone-salmeterol, Formoterol-mometasone Leukotriene modifiers: Montelukast, Zafirlukast, Zileuton Methylxanthines: Theophylline Short-acting, inhaled beta-2 agonists: Albuterol, Levalbuterol</p>
<p>Documentation</p>	<p>Claims based measure only. Link any medications used to treat asthma or other respiratory conditions with the applicable diagnosis.</p>
<p>Considerations</p>	<p>Educate patients about the difference between controller and reliever medication. Consider using 90-day prescriptions.</p>

<p>CODES:</p>	
<p>Acute Respiratory Failure ICD10</p>	<p>J96.00-J96.02, J96.20-J96.22</p>
<p>Asthma ICD10 SNOMED</p>	<p>J45.22, J45.30-J45.32, J45.40-J45.42, J45.50- J45.52, J45.901-J45.902, J45.909, J45.991, J45.998 There are over 100 SNOMED codes (one example is 11641008 Millers' asthma disorder)</p>
<p>Chronic Respiratory Conditions Due to Fumes/Vapors ICD10 SNOMED</p>	<p>J68.4, 506.4 15908004, 31803008, 32544004, 43098002, 61233003, 66110007, 69454006, 72163003, 74800004, 196025000, 196026004, 308905009</p>
<p>COPD ICD10 SNOMED</p>	<p>J44.0-J44.1, J44.9 13645005, 135836000, 195951007, 196001008, 285381006, 313296004, 313297008, 313299006, 1751000119100, 106001000119101</p>
<p>Cystic Fibrosis ICD10 SNOMED</p>	<p>E84.0, E84.11, E84.19, E84.8-E84.9 81423003, 86092005, 86555001, 190905008, 190909002, 235978006, 720401009, 762269004, 762270003, 762271004</p>
<p>ED ICD10 SNOMED</p>	<p>99281-99285 4525004</p>
<p>Emphysema ICD10 SNOMED</p>	<p>J43.0-J43.2, J43.8-J43.9 2912004, 4981000, 16003001, 16838000, 16846004, 23851004, 23958009, 31898008, 45145000, 47895001, 54288002, 57686001, 60805002, 68328006, 86680006, 87433001, 195957006, 195958001, 195959009, 195963002, 196026004, 233674008, 233675009, 233677001, 266355005, 266356006, 708030004</p>
<p>Obstructive Chronic Bronchitis ICD10 SNOMED</p>	<p>491.20-491.22 185086009, 2932410000119100</p>
<p>Other Emphysema ICD10 SNOMED</p>	<p>J98.2-J98.3 33325001, 77690003</p>

CHILD AND ADOLESCENT WELL-CARE VISITS (WCV)

<p>Measure Description</p> <p>Measure Source: HEDIS®</p>	<p>To assess the percentage of patients 3-21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.</p> <p>Three age stratifications are reported and total rate:</p> <ul style="list-style-type: none"> ■ 3-11 years old ■ 12-17 years old ■ 18-21 years old ■ Total
<p>Exclusions</p>	<p>Patients who use hospice services or choose to use hospice anytime in the measurement year.</p>
<p>Documentation</p>	<p>Claims based measure, meaning information is collected through claims data only to determine a rate. No chart reviews. (Note: this use to be a hybrid reported measure)</p>
<p>Considerations</p>	<p>The well-child visit must occur with a PCP, but the PCP does not have to be the practitioner assigned to the child.</p> <p>Visits can be via Telehealth.</p> <p>These are the previous measures: Well Child 3rd, 4th, 5th and 6th years of life combined with Adolescent Well Care Visits.</p> <p>Hybrid (medical record review) collection is no longer allowed.</p>
<p>CODES:</p>	
<p>Well Care</p>	<p>ICD10 Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z02.5, Z76.1, Z76.2</p> <p>CPT 99381,99382,99383,99384,99385,99391,99392,99393,99394,99395,99461</p> <p>HCPCS G0438, G0439, S0302</p> <p>SNOMED 10374000, 170099002, 170107008, 170114005, 170123008, 170132005, 170141000, 170150003, 170159002, 170168000, 170250008, 170254004, 170263002, 170272005, 170281004, 170290006, 170300004, 170309003, 171387006, 171394009, 171395005, 171409007, 171410002, 171416008, 171417004, 243788004, 268563000, 270356004, 401140000, 410620009, 410621008, 410622001, 410623006, 410624000, 410625004, 410626003, 410627007, 410628002, 410629005, 410630000, 410631001, 410632008, 410633003, 410634009, 410635005, 410636006, 410637002, 410638007, 410639004, 410640002, 410641003, 410642005, 410643000, 410644006, 410645007, 410646008, 410647004, 410648009, 410649001, 410650001, 442162000, 783260003, 444971000124105, 446301000124108, 446381000124104, 669251000168104, 669261000168102, 669271000168108, 669281000168106</p>

1 CPT® is a registered trademark of the American Medical Association. ICD-10-CM is the International Classification of Diseases, Tenth Revision, Clinical Modification. Source: MA 5 Star Technical Specifications and NCQA HEDIS® 2022; Technical Specifications for Health Plans HEDIS® measures and specifications are not a substitute for clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications. Documentation is for training and educational purposes.

<p>Measure Description</p> <p>Measure Source: HEDIS®</p>	<p>Patients 12 years of age and older screened for depression on the date of the encounter using an age appropriate standardized tool.</p> <p>All patients 12 years of age or older before the beginning of the measurement period with at least one eligible (any E/M office visit CPT code) encounter with a PCP during the measurement period are included.</p>
<p>Documentation</p>	<p>Depression screening during an eligible encounter based on claims data.</p>
<p>CODES:</p>	
<p>Depression Screening HCPCS</p>	<p>96127 Emotional & Behavioral Assessment Ages 12-18</p>

1 CPT® is a registered trademark of the American Medical Association. ICD-10-CM is the International Classification of Diseases, Tenth Revision, Clinical Modification. Source: MA 5 Star Technical Specifications and NCQA HEDIS® 2022; Technical Specifications for Health Plans HEDIS® measures and specifications are not a substitute for clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications. Documentation is for training and educational purposes.

WEIGHT ASSESSMENT AND COUNSELING FOR NUTRITION AND PHYSICAL ACTIVITY FOR CHILDREN/ ADOLESCENTS (WCC)


Measure Description		The percentage of patients 3–17 years of age who had an outpatient visit with a PCP or OB/GYN and who had evidence of the following during the measurement year. <ul style="list-style-type: none"> ■ BMI percentile documentation ■ Counseling for nutrition ■ Counseling for physical activity
Measure Source: HEDIS®		
Documentation Requirements	BMI	<p>Must include height, weight and BMI percentile during the measurement year. The height, weight and BMI percentile must be from the same data source. The following meets criteria for BMI percentile:</p> <ul style="list-style-type: none"> ■ BMI percentile documented as a value (e.g., 85th percentile). ■ BMI percentile plotted on an age-growth chart. <p>Patients' collected values (height, weight, BMI percentile) can be used. Ranges and thresholds do not meet the criteria and a distinct BMI % is required. Services rendered during a telephone visit, e-visit or virtual check-in meet criteria for BMI percentile indicator.</p>
	Nutrition Counseling	<p>Notes must include the date and at least one of the following:</p> <ul style="list-style-type: none"> ■ Discussion of current nutrition behaviors (e.g., eating habits, dieting behaviors) ■ Checklist indicating nutrition was addressed ■ Counseling or referral for nutrition education ■ Patient received educational materials on nutrition during a face-to-face visit ■ Anticipatory guidance for nutrition ■ Weight or obesity counseling
	Physical Activity	<p>Notes must include the date and at least one of the following:</p> <ul style="list-style-type: none"> ■ Discussion of current physical activity behaviors (e.g., exercise routine, participation in sports activities, exam for sports participation). ■ Checklist indicating physical activity was addressed. ■ Counseling or referral for physical activity. ■ Patient received educational materials on physical activity during a face-to-face visit. ■ Anticipatory guidance specific to the child's physical activity. ■ Weight or obesity counseling. <p>Exclusions:</p> <ul style="list-style-type: none"> ■ Patients who use hospice services or chose to use hospice anytime in measurement year.
CODES:		
BMI	ICD10	Z68.51-Z68.54
Nutrition Counseling	ICD10	Z71.3
	CPT	97802-97804
	HCPCS	G0270-G0271, G0447, S9449, S9452, S9470
	SNOMED	There are over 65 Snomed codes; one example is 11816003 Diet education (procedure)
Physical Activity	ICD10	Z02.5, Z71.82
	HCPCS	G0447, S9451
	SNOMED	103736005, 171356009, 171357000, 171358005, 171359002, 171360007, 171361006, 183073003, 183075005, 223415003, 223440005, 281090004, 304517008, 304549008, 304558001, 310882002, 386291006, 386292004, 386463000, 390864007, 390893007, 398636004, 398752005, 408289007, 410200000, 410289001, 410335001, 426866005, 429095004, 429778002, 710849009, 435551000124105

<p>Measure Description</p> <p>Measure Source: HEDIS®</p>	<p>The percentage of members who had the following number of well-child visits with a PCP during the last 15 months. The following rates are reported:</p> <ol style="list-style-type: none"> 1. Well-Child Visits in the First 15 Months. Children who turned 15 months old during the measurement year: Six or more well-child visits. 2. Well-Child Visits for Age 15 Months–30 Months. Children who turned 30 months old during the measurement year: Two or more well-child visits.
<p>Exclusions</p>	<p>Patients who use hospice services or chose to use hospice anytime in measurement year</p>
<p>Documentation</p>	<p>Claims based measure, meaning information is collected through claims data only to determine a rate. No chart reviews. (Note: this use to be a hybrid reported measure)</p>
<p>Considerations</p>	<p>The well-child visit must occur with a PCP, but the PCP does not have to be the practitioner assigned to the child.</p> <p>Visits can be via Telehealth.</p>
<p>CODES:</p>	
<p>Well Care Visit</p>	<p>ICD10 Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z02.5, Z76.1, Z76.2</p> <p>CPT 99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461</p> <p>HCPCS G0438, G0439, S0302</p> <p>SNOMED 170099002, 170107008, 170114005, 170123008, 170132005, 170141000, 170150003, 170159002, 170168000, 170250008, 170254004, 170263002, 170272005, 170281004, 170290006, 170300004, 170309003, 171387006, 171394009, 171395005, 171409007, 171410002, 171416008, 171417004, 243788004, 268563000, 270356004, 401140000, 410620009, 410621008, 410622001, 410623006, 410624000, 410625004, 410626003, 410627007, 410628002, 410629005, 410630000, 410631001, 410632008, 410633003, 410634009, 410635005, 410636006, 410637002, 410638007, 410639004, 410640002, 410641003, 410642005, 410643000, 410644006, 410645007, 410646008, 410647004, 410648009, 410649001, 410650001, 442162000, 783260003, 444971000124105, 446301000124108, 446381000124104, 669251000168104, 669261000168102, 669271000168108, 669281000168106</p>

1 CPT® is a registered trademark of the American Medical Association. ICD-10-CM is the International Classification of Diseases, Tenth Revision, Clinical Modification. Source: MA 5 Star Technical Specifications and NCQA HEDIS® 2022; Technical Specifications for Health Plans HEDIS® measures and specifications are not a substitute for clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications. Documentation is for training and educational purposes.



*KEEP GETTING BETTER



Hackensack Meridian
Partners

HackensackMeridianHealthPartners.org

HMH-4762-05.22-LD